



Banks, A. M., Barker, G. L. A., Bailey, A. M., & Foster, G. D. (2017). Draft Genome Sequence of the Coprinoid Mushroom *Coprinopsis strossmayeri*. *Genome Announcements*, 5(14), [e0004417].  
<https://doi.org/10.1128/genomeA.00044-17>

Publisher's PDF, also known as Version of record

License (if available):  
CC BY

Link to published version (if available):  
[10.1128/genomeA.00044-17](https://doi.org/10.1128/genomeA.00044-17)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the final published version of the article (version of record). It first appeared online via American Society for Microbiology at <https://doi.org/10.1128/genomeA.00044-17> . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>



# Draft Genome Sequence of the Coprinoïd Mushroom *Coprinopsis strossmayeri*

Alice M. Banks,\* Gary L. A. Barker, Andy M. Bailey, Gary D. Foster

School of Biological Sciences, University of Bristol, Bristol, United Kingdom

**ABSTRACT** *Coprinopsis strossmayeri* is a coprinoid mushroom favoring the habitat of herbivore dung. As a result of this highly competitive environment, *C. strossmayeri* is anticipated to produce a wide array of antimicrobial secondary metabolites (SMs) of potential pharmaceutical importance. Here, we present the draft genome sequence of *C. strossmayeri*.

Basidiomycete fungi are part of a hugely varied phylum of organisms capable of carrying out a diversity of important roles (1). The coprinoid mushrooms encompass fungi from the *Coprinus*, *Coprinopsis*, and *Coprinellus* genera, and, along with the *Parasola*, comprise the Psathyrellaceae family (2). Most coprinoids favor a habitat of herbivore dung, but growth on decaying woodland material is also common (3). A likely result of residing in such competitive environments and cohabiting with predatory microorganisms is the production of a plethora of bioactive secondary metabolites (SMs) used in defense, many of which can potentially be exploited for human use (4). Terpenoids are the most abundant class of compounds produced by basidiomycetes (5), the derivatives of which have been developed for medical applications, such as the diterpene antibiotic pleuromutilin from *Clitopilus passeckerianus* (6).

The strain sequenced was obtained from the CBS Fungal Biodiversity Centre, submitted as *Coprinus quadrididus* CBS 177.39. The internal transcribed spacer region was amplified and sequenced to confirm species identity. Subsequent analysis by BLASTn search identified the isolate as *Coprinopsis strossmayeri*, showing 98% identity to *C. strossmayeri* strain SZMC-NL-0774 (GenBank accession number HQ847048.1).

The genomic DNA of dikaryotic *C. strossmayeri* (CBS 177.39) mycelium was sequenced using the Illumina HiSeq 2500 system. Paired-end 100-bp fragments were prepared and sequenced, generating 93,872,638 reads covering a total 9,387 Mbp ( $\geq$ Q30 bases, 91.87%). Data were processed using RTA version 1.17.21.3, with default settings, and reads were demultiplexed, allowing no mismatches, with CASAVA 1.8.2. Quality-trimmed reads were assembled using CLC Genomics Workbench 6. This resulted in a genome assembly comprising 622 contigs totaling 33,316,483 bp, with an average contig length of 53,563 bp (largest contig, 1,580,550 bp; smallest contig, 1,000 bp). An  $N_{25}$  contig length of 380,611 bp,  $N_{50}$  of 190,582 bp, and  $N_{75}$  of 80,746 bp were obtained. The G+C content was 49%. This genome size is comparable to those of related coprinoid fungi.

Genome analysis was performed using antiSMASH (7). This located five terpene synthase genes related to terpenoid biosynthesis, four of which are characteristic of sesquiterpene biosynthesis, while the other is expected to be involved with diterpene biosynthesis, based on phylogenetic analyses. A local BLAST search of the genome located two geranylgeranyl diphosphate synthases likely to be involved with diterpene biosynthesis. Terpenoid production is well documented among coprinoid fungi; many products of this chemical class have been reported from related species (8–11). One type I partially reducing polyketide synthase and one nonribosomal peptide synthase

**Received** 20 January 2017 **Accepted** 7 February 2017 **Published** 6 April 2017

**Citation** Banks AM, Barker GLA, Bailey AM, Foster GD. 2017. Draft genome sequence of the coprinoid mushroom *Coprinopsis strossmayeri*. Genome Announc 5:e00044-17. <https://doi.org/10.1128/genomeA.00044-17>.

**Copyright** © 2017 Banks et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Gary D. Foster, [Gary.Foster@bristol.ac.uk](mailto:Gary.Foster@bristol.ac.uk).

\* Present address: Alice M. Banks, School of Biology, Newcastle University, Newcastle upon Tyne, United Kingdom.

were also identified through antiSMASH. Interestingly, a locus showing similarity to copsisin, an antimicrobial peptide from *Coprinopsis cinerea*, was located through a BLAST search (12). We plan to identify the biosynthetic gene clusters in *C. strossmayeri* responsible for the production of novel antimicrobial SMs and to characterize the genes constituting these pathways. The numbers of genes and gene clusters identified highlight the potential of *C. strossmayeri* as a source of novel terpenoid SMs to be exploited in the pharmaceutical industry.

**Accession number(s).** This whole-genome shotgun project has been deposited in DDBJ/ENA/GenBank under the accession numbers [FTPT01000001](#) to [FTPT01000622](#).

## ACKNOWLEDGMENT

We acknowledge the CBS Fungal Biodiversity Centre, Utrecht, Netherlands for access to the fungal stock collection.

## REFERENCES

- de Mattos-Shipley KMJ, Ford KL, Alberti F, Banks AM, Bailey AM, Foster GD. 2016. The good, the bad and the tasty: the many roles of mushrooms. *Stud Mycol* 85:125–157. <https://doi.org/10.1016/j.simyco.2016.11.002>.
- Redhead SA, Vilgalys R, Moncalvo J-M, Johnson J, Hopple JS. 2001. *Coprinus* Pers. and the disposition of *Coprinus* species *sensu lato*. *Taxon* 50:203. <https://doi.org/10.2307/1224525>.
- Badalyan SM, Szafranski K, Hoegger PJ, Navarro-González M, Majcherzyk A, Kües U. 2011. New Armenian wood-associated coprinoid mushrooms: *Coprinopsis strossmayeri* and *Coprinellus* aff. *radians*. *Diversity* 3:136–154. <https://doi.org/10.3390/d3010136>.
- Keller NP, Turner G, Bennett JW. 2005. Fungal secondary metabolism—from biochemistry to genomics. *Nat Rev Microbiol* 3:937–947. <https://doi.org/10.1038/nrmicro1286>.
- Abraham WR. 2001. Bioactive sesquiterpenes produced by fungi: are they useful for humans as well? *Curr Med Chem* 8:583–606. <https://doi.org/10.2174/0929867013373147>.
- Bailey AM, Alberti F, Kilaru S, Collins CM, de Mattos-Shipley K, Hartley AJ, Hayes P, Griffin A, Lazarus CM, Cox RJ, Willis CL, O'Dwyer K, Spence DW, Foster GD. 2016. Identification and manipulation of the pleuromutilin gene cluster from *Clitopilus passeckerianus* for increased rapid antibiotic production. *Sci Rep* 6:25202. <https://doi.org/10.1038/srep25202>.
- Weber T, Blin K, Duddela S, Krug D, Kim HU, Bruccoleri R, Lee SY, Fischbach MA, Müller R, Wohlleben W, Breitling R, Takano E, Medema MH. 2015. antiSMASH 3.0—a comprehensive resource for the genome mining of biosynthetic gene clusters. *Nucleic Acids Res* 43:W237–W243. <https://doi.org/10.1093/nar/gkv437>.
- Badalyan SM, Kües U, Melikyan LR, Navarro-González M. 2005. Medicinal properties of coprinoid mushrooms (Basidiomycetes, Agaricales). *Int J Med Mushrooms* 7:378–380.
- Reina M, Orihuela JC, González-Coloma A, de Inés C, de la Cruz M, González del Val A, Torno JR, Fraga BM. 2004. Four illudane sesquiterpenes from *Coprinopsis episcopalis*. *Phytochemistry* 65:381–385. <https://doi.org/10.1016/j.phytochem.2003.10.023>.
- Gonzalez del Val A, Platas G, Arenal F, Orihuela JC, Garcia M, Hernández P, Royo I, De Pedro N, Silver LL, Young K, Vicente MF, Pelaez F. 2003. Novel illudins from *Coprinopsis episcopalis* (syn. *Coprinus episcopalis*), and the distribution of illudin-like compounds among filamentous fungi. *Mycol Res* 107:1201–1209. <https://doi.org/10.1017/S0953756203008487>.
- Kettering M, Valdivia C, Sterner O, Anke H, Thines E. 2005. Heptemerones A–G, seven novel diterpenoids from *Coprinus heptemerus*: producing organism, fermentation, isolation and biological activities. *J Antibiot (Tokyo)* 58:390–396. <https://doi.org/10.1038/ja.2005.49>.
- Essig A, Hofmann D, Münch D, Gayathri S, Künzler M, Kallio PT, Sahl HG, Wider G, Schneider T, Aebi M. 2014. Copsisin, a novel peptide-based fungal antibiotic interfering with the peptidoglycan synthesis. *J Biol Chem* 289:34953–34964. <https://doi.org/10.1074/jbc.M114.599878>.